

Chemistry and kinematics of the pre-stellar core L1544: Constraints from H_2D^+ .

Floris van der Tak¹, Paola Caselli², Malcolm Walmsley², Cecilia Ceccarelli³, Aurore Bacmann⁴, and Antonio Crapsi²

¹ Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53111 Bonn, Germany vdtak@mpifr-bonn.mpg.de

² Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125 Firenze, Italy

³ Laboratoire d'Astrophysique de l'Observatoire de Grenoble, BP 53, 38041 Grenoble Cedex, France

⁴ European Southern Observatory, Karl-Schwarzschild-Straße 2, 85748 Garching bei München, Germany

This paper explores the sensitivity of line profiles of H_2D^+ , HCO^+ and N_2H^+ , observed towards the center of L 1544, to various kinematic and chemical parameters. The total width of the H_2D^+ line can be matched by a static model and by models invoking ambipolar diffusion and gravitational collapse. The derived turbulent line width is $b=0.15 \text{ km s}^{-1}$ for the static case and $\lesssim 0.05 \text{ km s}^{-1}$ for the collapse case. However, line profiles of HC^{18}O^+ and N_2H^+ rule out the static solution. The double-peaked H_2D^+ line shape requires either infall speeds in the center that are much higher than predicted by ambipolar diffusion models, or a shell-type distribution of H_2D^+ , as is the case for HCO^+ and N_2H^+ . At an offset of $\approx 20''$ from the dust peak, the H_2D^+ abundance drops by a factor of ≈ 5 .

1 Introduction

Deuterium-bearing molecules are important as probes of the very cold phases of molecular clouds prior to star formation. The H_2D^+ ion is especially important as tracer of H_3^+ , the primary ion in dense molecular clouds, which does not have a dipole moment and hence no pure rotational lines. In addition, at low temperatures ($\lesssim 10 \text{ K}$), H_2D^+ has the ability to channel D atoms from their main reservoir, HD, into heavier species. This process leads to abundance ratios of $\text{DCO}^+/\text{HCO}^+$ and $\text{N}_2\text{D}^+/\text{N}_2\text{H}^+$ of $\sim 10^{-3} - 10^{-1}$ observed towards dense cores, much larger than the elemental D/H ratio of $\sim 10^{-5}$. Recent observations of multiply deuterated H_2CO , CH_3OH , H_2S and NH_3 (see Ceccarelli, this volume) suggest that under extreme conditions, a significant fraction of D may be transferred to heavy molecules. We wish to quantify the role of H_2D^+ in this process, and compare with other processes such as grain surface reactions (Caselli, this volume).

The ground-state $1_{01}-0_{00}$ transition of para- H_2D^+ at 1370 GHz will be a prime target for GREAT on SOFIA. However, with the upper energy level

65 K above ground, this line will only be excited in relatively warm ($\gtrsim 20$ K) regions, where chemical fractionation is ineffective. For colder sources, the $1_{10}-1_{11}$ ground-state transition of ortho- H_2D^+ at 372 GHz is more suitable, which can be observed from the ground under good conditions. At the low temperatures ($\lesssim 10$ K) and high densities ($> 10^5 \text{ cm}^{-3}$) of pre-stellar cores, reactive collisions with ortho- H_2 keep the ortho-para ratio of H_2D^+ at ~ 1 , orders of magnitude above the LTE value [1, 2].

Until 2002, only two detections of H_2D^+ had been obtained, which indicated abundances of $\sim 10^{-11}$ – 10^{-12} towards Class 0 objects [3, 4]. In October 2002, we [5] observed strong H_2D^+ emission towards the pre-stellar core L 1544, and derived an abundance of $\sim 1 \times 10^{-9}$ in the central $\approx 20''$. Such a high abundance suggests that in this region, all CNO-bearing species are depleted onto dust grains, a situation explored in more detail by Walmsley (this volume). Data taken in June 2003 at the CSO indicate that the same phenomenon takes place in at least five other pre-stellar cores. More observations are scheduled for December 2003. These data will be presented in a forthcoming paper. Here we investigate the line profile of H_2D^+ in L 1544, and derive its abundance outside the central region.

2 Kinematics

The line profile of H_2D^+ towards L 1544 appears double-peaked, although the signal-to-noise ratio is not high (Fig. 1). Fitting two Gaussians to the profile yields results very similar to those for the profiles of HC^{18}O^+ , D^{13}CO^+ and N_2H^+ , observed by [6]: two thermal components separated by $\approx 0.26 \text{ km s}^{-1}$ (Table 1). We therefore investigate whether the kinematic models that fit the HCO^+ and N_2H^+ data also reproduce the H_2D^+ line profile.

Table 1. Centroids and widths of the two velocity components. Numbers in brackets denote uncertainties in units of the last decimal.

Line	V_{LSR} km s^{-1}	ΔV_{obs} km s^{-1}	ΔV_{T}^a km s^{-1}
$\text{H}_2\text{D}^+ (1_{10}-1_{11})$	7.06(3) 7.34(2)	0.22(5) 0.25(6)	0.28–0.34
$\text{HC}^{18}\text{O}^+ (1-0)$	7.04(1) 7.28(1)	0.18(3) 0.23(3)	0.10–0.12
$\text{D}^{13}\text{CO}^+ (2-1)$	7.08(2) 7.35(4)	0.20(4) 0.20(8)	0.10–0.12
$\text{N}_2\text{H}^+ (1-0, F_1 F=10-11)$	7.08(1) 7.33(1)	0.19(1) 0.20(2)	0.11–0.13

^a Thermal line width at $T_{\text{kin}}=7$ and 10 K.




Fig. 1. Observed line profiles of H_2D^+ (left), HC^{18}O^+ (middle) and N_2H^+ (right), and synthetic profiles for the static (dotted) and infall (dashed) models.

The line profile of H_2D^+ was modeled using a Monte Carlo radiative transfer program [7]⁵. Figure 2 shows the adopted temperature and density structure of L1544, taken from [8]. See [5] for details of the excitation model; at $T_{\text{kin}} \lesssim 10$ K, ortho- H_2D^+ is essentially a two-level system, so that our results are not sensitive to the collision rates of non-radiative transitions between high-lying levels. For H_2D^+ we adopt an abundance of 1×10^{-9} in the central $20''$ [5]. For HCO^+ , DCO^+ , N_2H^+ and N_2D^+ , we used the abundance profiles from [9], Model 3, and assumed zero abundance inside $r=2500$ AU. We explored static models, and models with velocity fields from the ambipolar diffusion models ‘t3’ and ‘t5’ of [10] (see [6] for details). For the turbulent broadening, Doppler parameters b between 0.05 and 0.25 km s^{-1} were tried. Smaller values of b are overwhelmed by thermal broadening; larger ones do not fit the data.

We find that the total width of the H_2D^+ line can be matched using either velocity field. While the best-fit static model has $b=0.15$, $b=0.05$ gives the best fits with the infall velocity fields. However, the HC^{18}O^+ and N_2H^+ $J=1-0$ observations from [6] rule out the static model, which does not give a double-peaked line shape. Using the infall velocity fields, the data are matched with $b=0.05$, consistent with the H_2D^+ results (Fig. 1).

None of our adopted velocity fields reproduces the double-peaked H_2D^+ line shape that the observations indicate. One possibility is that the infall speeds of ≈ 0.1 km s^{-1} continue further inwards than in the models by [10]. Alternatively, the distribution of H_2D^+ may have a central hole, not because of adsorption onto dust grains (as for CO and N_2 , the precursors of HCO^+ and N_2H^+), but due to conversion into D_2H^+ and D_3^+ .

⁵ <http://www.mpifr-bonn.mpg.de/staff/fvandertak/ratran/>




Fig. 2. Radial profiles of dust and gas temperature, and the densities of H_2 , H_2D^+ ($\times 2 \times 10^9$), HCO^+ ($\times 10^{12}$) and N_2H^+ ($\times 10^{10}$) in our model of L 1544.

3 Abundance profile of H_2D^+

The abundance of H_2D^+ away from the dust peak of L 1544 was estimated to be a factor of two lower than toward the dust peak [5]. However, this abundance may be an overestimate because some fraction of the emission at the $20''$ offset positions is pickup from the central core. We have run Monte Carlo models of the H_2D^+ emission, using the same temperature and density structure as before, and dropping the H_2D^+ abundance at a $20''$ radius by factors of 2–10 from its central value of 1×10^{-9} . The H_2D^+ intensity at the $20''$ offset position is best matched if the abundance drops by a factor of ≈ 5 at this radius. This result is independent of the velocity field. Models where this factor is 3 or 10 produce clearly worse matches to the data.

In a future paper, we will follow our results up with two-dimensional models and an exploration of different velocity fields. We will also model line profiles at offset positions.

References

1. Pagani, L., Salez, M., & Wannier, P. G.: A&A, 258, 479 (1992)
2. Gerlich, D., Herbst, E., & Roueff, E.: Plan. Sp. Science, 50, 1275 (2002)
3. Stark, R., van der Tak, F. F. S., & van Dishoeck, E. F.: ApJ, 521, L67 (1999)
4. Stark, R., Sandell, G., Beck, S., et al.: ApJ, submitted (2003)
5. Caselli, P., van der Tak, F. F. S., Ceccarelli, C., & Bacmann, A.: A&A, 403, L37 (2003)
6. Caselli, P., Walmsley, C. M., Zucconi, A., Tafalla, M., Dore, L., & Myers, P. C.: ApJ, 565, 331 (2002)
7. Hogerheijde, M.R., & van der Tak, F. F. S.: A&A, 362, 697 (2000)
8. Galli, D., Walmsley, M., & Gonçalves, J.: A&A, 394, 275 (2002)
9. Caselli, P., Walmsley, C. M., Zucconi, A., Tafalla, M., Dore, L., & Myers, P. C.: ApJ, 565, 344 (2002)
10. Ciolek, G. E. & Basu, S.: ApJ, 529, 925 (2000)